

/

(Prunus persica)

(2) (1)
(4) (3)

63 126 .

Meloidogyne incognita

%44 (Kofoid and White, 1919) Chitwood 1949
330-20 3 100 / 90-10
10

3 100 / 36 30 %23.1 %4 *Pratylenchus* spp.

Xiphinema spp.

3 100 / 28 %11

X. intermedium Lamberti *X. pyrenaicum* Dalmasso, 1964

Paratylenchus spp. & Bleve-Zacheo, 1979

Tylenchus *Helicotylenchus* spp. *Amplimerlinius* spp.

46 52 223 %14 %21 %36 %26 spp.
3 100 / 29

Amplimerlinius spp. *Paratylenchus* spp.

%38.5 %3.8 %11.5 *Helicotylenchus* spp.
3 100 120 29 18

M. incognita, Pratylenchus, Xiphinema, Prunus, :

30621 . .

(1)
(2)
(4) (3)

/

First Report: Plant Parasitic Nematodes Associated with Peachs (*Prunus persica*) at Damascus and Hama Governorates, Syria

M. Al-Qasim⁽¹⁾; K. Assas⁽²⁾;
L. Banna⁽³⁾ and W. Abu-Gharbieh⁽⁴⁾

ABSTRACT

A study was conducted to investigate the presence of root knot nematode (RKN) and other plant parasitic nematodes associated with Syrian most important peach plantation areas. 126 soil and root samples were taken from 63 peach orchards of Ghuta, Zabadani, Yabrod, Syrghaya and Kharabo at Damascus governorate, and Tezeen, Kufr Ton, Khattab, Arzeh, Sheha, Qamhaneh, Balhasen, Rastan, Mhardeh, Salmieh, Sqalbieh and Sharaya at Hama governorate. We are reporting that *Meloidogyne incognita* (Kofoid and White, 1919) Chitwood 1949 infects 44% of orchards with an average intensity range of 10-90 second stage juveniles per 100 cm³ and average severity range of 20-330 galls per 10 gm roots. RKN did not noticed at any of Damascus governorate peach orchards, however, peach transplants at Al-Nasrieh nursery were severely infected with RKN and problem was solved by eliminating all transplants. *Pratylenchus* spp. presents in 4% and 23.1% of Hama and Damascus governorates with an average intensity of 30 and 36 nematodes per 100 cm³ soil, respectively. *Xiphinema* spp. presented in 11% of sampled Hama orchards with an average intensity of 28 nematodes per 100 cm³ and was not noticed at Damascus governorate. The species; *X. pyrenaicum* Dalmasso, 1964 and *X. intermedium* Lamberti & Bleve-Zacheo, 1979 were classified. *Paratylenchus* spp., *Amplimerlinius* spp., *Helicotylenchus* spp. and *Tylenchus* spp. presents in 26%, 36%, 21%, and 14% of sampled Hama orchards with an average intensity of 223, 52, 46 and 29 nematodes per 100 cm³ soil, respectively. *Paratylenchus* spp., *Amplimerlinius* spp. and *Helicotylenchus* spp. were presented in 11.5%, 3.8 and 23.5 of sampled Damascus orchards, with an average intensity of 18, 26 and 120 nematodes per 100 cm³. This is the first report of plant parasitic nematode associated with Syrian peach orchards.

Key words: *M. incognita*, Peach, *Prunus*, *Pratylenchus*, *Xiphinema*

⁽¹⁾ National Center for Agricultural Research and Extension, MOA, Amman, Jordan.

⁽²⁾ Plant Protection Department, Faculty of Agriculture, Damascus University, P.O. Box 30621, Damascus, Syria.

^{(3),(4)} Plant Protection Department, Faculty of Agriculture, University of Jordan, Amman, Jordan..

50.1
2006
%13.9 2006)
2006 65.9 1997
20.6 23.0
% 69 35.7
(2006) 2006

Sasser and Freckman,) %12.3

.(1987

(Ogawa *et al.*, 1995)

.(2003)

(Lamberti, 1984)

Ogawa *et al.*,)

(1995

Meloidogyne incognita(Kofoid and White, 1919) Chitwood 1949; *M. javanica* (Treub, 1885)Chitwood, 1949; *M. arenaria* (Neal, 1889) Chitwood *et al.*, 1952*P. vulnus* Allen and Jensen, 1951; *Pratylenchus zeae*Graham, 1951; *P. crenatus* Sher & Allen's, 1953; *P. brachyurus**C. xenoplax* (Raski, 1952) Godfrey, 1929; *P. penetrans*Luc & Raski, 1981, *Criconemella curvata* (Raski, 1952) Luc & Raski,

1981

Xiphinema americanum Cobb 1913

Peach Rosette Mosaic Virus (PRMV)

;Ogawa, 1995) Peach Yellow Bud Mosaic (PYBMV)

.(2002 ; Barker and Clayton, 1973; Whitehead, 1998

Nyczepir) *M. hapla* Chitwood, 1949*Pratylenchus crenatus* Loof, (and Halbrecht, 1993*P. thornei* Sher and Allen, 1953 (Norton *et al.*, 1984) 1960Siddiqui) *P. neglectus* (Rensch, 1924) Filipjev & Schur. Stet., 1941*P. penetrans* (Cobb, 1917) Chitwood & Oteifa, 1952 (*et al.*, 1973*Criconemella* .(Barker and Clayton, 1973)*C. simile* (Cobb, 1918) *curvata* (Raski, 1952) Luc & Raski, 1981*C. xenoplax* (Raski, 1952) Luc & Raski, 1981 1922 Chitwood, 1949.(Ogawa, 1995; Jaffee *et al.* 1987)*Longidorus elongates* (De Man, 1876) Thorne & Swanger, 1936*Xiphinema* .(Knight *et al.*, 1997)*americanum* Cobb, 1913

.(Meyer and Hugo, 1994)

Xiphinema pachtaicum (Tulaganov,1938) Kirjanova, 1951

Helicotylenchus

Prunus amygdalus *varicaudatus* Yuen, 1964

H. dihystra (Cobb, 1893) Sher, 1961 .Batsch

.(Lamberti, 1984) *Prunus domestica* L.

63

126

30 - 15

.(Barker *et al.*, 1985)

.(Barker *et al.*, 1985)

5

Baermann Funnel Cobb Sieving

³ 100

(Barker *et al.*, 1985)

40

48

5

(%4)

5

.(Jacob and van Bezooijen, 1984) %2

Grooved

.(x45)

slide

Seinhorst,)

(1959)

10

10

72

2-1

10

(x45)

Perennial Patterns
(Barker *et al.*, 1985)

(

)

Soil

Soil EC

Soil pH

Texture

(1995)

³

10 /

10 /
330

330-20

(

125 315)

10 /

113

%44

90 -10

(1)

³

100 /

Meloidogyne incognita (Kofoid and White, 1919) Chitwood 1949

** * (1)

Meloidogyne incognita

	()	()	(J2)								
				EC	pH	%	%	%			
	8 (0)	0 %	-	-	-	-	-	-	-	-	-
	6 (0)	0 %	-	-	-	-	-	-	-	-	-
	10 (4)	40 %	15-40 (25)	10-50 (30)	1.55	7.8	57.5	12.5	30	SCL	
	6 (4)	67 %	180-450 (315)	50-150 (90)	1.5	7.6	72	13.5	14.5	SL	
	12 (10)	83 %	10-80 (36)	10-70 (30)	1.62	7.9	60	15	25	SCL	
	12 (8)	67 %	10-30 (20)	10-50 (30)	2.1	7.5	53	17	30	SCL	
	12 (8)	67 %	10-60 (35)	20-60 (50)	3.6	7.9	30	32.5	37.5	CL	
	6 (2)	33 %	270-390 (330)	20-40 (30)	1.45	7.7	75	12.5	12.5	SL	
	6 (6)	100%	10-60 (28)	0-20 (10)	1.7	7.7	52.5	17.5	30	SCL	
	6 (2)	33 %	10-240 (125)	0-20 (10)	1.8	8	75	12.5	12.5	SL	
	8 (0)	0 %	-	-	-	-	-	-	-	-	
	8 (0)	0 %	-	-	-	-	-	-	-	-	
Total/ Average	100 (44)	44 %	(113)	(35)	/						
	4 (0)	0 %	-	-	-	-	-	-	-	-	
	8 (0)	0 %	-	-	-	-	-	-	-	-	
	6 (0)	0 %	-	-	-	-	-	-	-	-	
	6 (0)	0 %	-	-	-	-	-	-	-	-	
	(0)2	0 %	-	-	-	-	-	-	-	-	
Total/ Average	(0)26	-	-	-	/						

×(/): (Norton, 1978) % 100 *
 10/ **
 . 3 100/J2

%4 *Pratylenchus*
 / 30
 %23.1 (2) 3 100
 . 3 100 / 36
Pratylenchus spp. ** * (2)
Xiphinema spp.

		<i>Pratylenchus</i> spp.			<i>Xiphinema</i> spp.		
		()		()	()		()
		8 (0)	0 %	-	8 (0)	0	-
		6 (0)	0 %	-	6 (0)	0	-
		10 (0)	0 %	-	10 (0)	0	-
		6 (0)	0 %	-	6 (2)	33 %	20-40 (30)
		12 (0)	0 %	-	12 (3)	25 %	10-50 (30)
		12 (0)	0 %	-	12 (0)	0	-
		12 (4)	33 %	10-60 (30)	12 (4)	33 %	10-60 (30)
		6 (0)	0 %	-	6 (2)	33 %	20 (20)
		6 (0)	0 %	-	6 (0)	0	-
		6 (0)	0 %	-	6 (0)	0	-
		8 (0)	0 %	-	8 (0)	0	-
		8 (0)	0 %	-	8 (0)	0	-
Total / Average		100 (4)	4.0 %	30	100 (11)	11.0 %	28
		4 (2)	50 %	40-80 (60)	4 (0)	0 %	-
		8 (0)	0 %	-	8 (0)	0 %	-
		6 (3)	50 %	10-40 (30)	6 (0)	0 %	-
		6 (0)	0 %	-	6 (0)	0 %	-
		(1)2	%50	18	2 (0)	0 %	-
Total / Average		26 (6)	23.1 %	36	26 (0)	0.0 %	-

.(Norton, 1978) %100×(/) *
 . 3 100 / **

%11 *Xiphinema* spp.
 28 (2)
 . 3 100 /

- *X. pyrenaicum* Dalmasso, 1964 (Dalmasso, 1969)
- *X. intermedium* Lamberti & Bleve-Zacheo, 1979 (Lamberti and Bleve-Zacheo, 1979)

$\frac{\%26}{100} \times \frac{223}{1035} = 0.0115$
 $\times 100 = 1.15\%$

$\frac{3}{100} \times 18 = 0.54$
 $\times 100 = 54\%$

Paratylenchus spp. ** * (3)
Amplimerlinius spp.

	<i>Paratylenchus</i>			<i>Amplimerlinius</i>		
	()	()	()	()	()	()
	8 (4)	50 %	20-50 (30)	8 (0)	0 %	-
	6 (2)	33 %	80-100 (90)	6 (0)	0 %	-
	10 (0)	0 %	-	10 (2)	20 %	20-40 (30)
	6 (0)	0 %	-	6 (2)	33 %	10-50 (30)
	12 (0)	0 %	-	12 (6)	50 %	30-120 (70)
	12 (0)	0 %	-	12 (0)	0 %	-
	12 (8)	67 %	60-200 (125)	12 (8)	67 %	80-220 (145)
	6 (4)	67 %	30-2040 (1035)	6 (2)	33 %	80-120 (100)
	6 (4)	67 %	10-60 (30)	6 (4)	67 %	10-80 (30)
	6 (0)	0 %	-	6 (0)	0 %	-
	8 (0)	0 %	-	8 (8)	100 %	10-60 (25)
	8 (4)	50 %	10-60 (30)	8 (4)	50 %	10-70 (30)
Total/Average	100 (26)	26.0 %	223	100 (36)	36.0 %	52
	4 (2)	50 %	30 (30)	4 (0)	0 %	-
	8 (0)	0 %	-	8 (0)	0 %	-
	6 (0)	0 %	-	6 (0)	0 %	-
	6 (0)	0 %	-	6 (0)	0 %	-
	2 (1)	50 %	7	2 (1)	50 %	29
Total/Average	26 (3)	11.5 %	18	26 (1)	3.8 %	29

$\frac{1}{100} \times 100 = 1\%$ *
 $\frac{3}{100} \times 100 = 3\%$ **

%36 *Amlimerlinius* spp.
Tylenchus spp. *Helicotylenchus* spp.
 3 100 / 29 46 52 %14 %21

%11.5 *Helicotylenchus* spp. *Paratylenchus* spp.
 %38.5
 3 100 / 120 18
 %3.8 *Amlimerlinius* spp.
 3 100 29
Tylenchus spp.
 .(4)

Tylenchus Helicotylenchus spp. (4) spp.

		<i>Helicotylenchus</i>			<i>Tylenchus</i>		
		()		()	()		()
		8 (4)	50 %	10-60 (30)	8 (0)	0 %	-
		6 (0)	0 %	-	6 (0)	0 %	-
		10 (0)	0 %	-	10 (0)	0 %	-
		6 (0)	0 %	-	6 (0)	0 %	-
		12 (3)	25 %	30-90 (50)	12 (2)	17 %	30 (30)
		12 (8)	67 %	10-80 (30)	12 (0)	0 %	-
		12 (8)	67 %	10-120 (50)	12 (10)	83 %	10-50 (27)
		6 (2)	33 %	20-100 (60)	6 (2)	33 %	20-40 (30)
		6 (0)	0 %	-	6 (0)	0 %	-
		6 (0)	0 %	-	6 (0)	0 %	-
		8 (0)	0 %	-	8 (0)	0 %	-
		8 (0)	0 %	-	8 (0)	0 %	-
Total /Average		100 (21)	21.0 %	46	100 (14)	14.0 %	29
		4 (0)	-	-	4 (0)	0 %	-
		8 (4)	50 %	10-70 (30)	8 (0)	0 %	-
		6 (3)	50 %	20-160 (90)	6 (0)	0 %	-
		6 (3)	50 %	120-380 (240)	6 (0)	0 %	-
		2 (0)	0	-	2 (0)	0 %	-
Total /Average		26 (10)	38.5 %	120	26 (0)	0.0 %	-

.(Norton, 1978) %100×(/) : *
 3 100 / **

Meloidogyne incognita

(Sasser and Carter, 1985)

(Lamberti, 1984; Al Ahmed, 1987)

(2003)

10 (%12.5 10/ 330 %12.5 %75.0:)
10/ (%37.5 %32.5 %30.0:)

(Stirling, 1991)

()

)

(

Nyczepir and)

(2003

2002

1995

Halbrendt, 1993

(
(1985) Sasser Carter

%16
(McKenry, 1989)

(2002)
Xiphinema spp.

%10
(Decraemer and Geraet, 2006) Michigan
Helicotylenchus spp. Steiner, 1945
Tylenchorhynchus spp. Cobb, 1913, Micoletzky, 1922
(Decraemer and Geraet, 2006)

REFERENCES

- .(2002) .
343 .
- .(1995) .
246 .
- .(2003) .
360 .
- .(1995) .
255 .
- .(2006) .
- Al-Ahmed, M. (1987). The status of plant- parasitic nematodes in cereals and food and forage legumes in Syria. In: Nematodes Parasitic to Cereals and Legumes in Temperate Semi-Arid Regions. M. C. Saxena, R. A. Sikora and J. P. Srivastava (Editors). ICARDA-135.
- Barker, K., and C. Clayton. (1973). Nematodes attacking cultivars of peach in North Carolina. *Journal of Nematology* 5:265-271.
- Barker, K., J. Sasser and C. Carter. (1985). An Advanced Treatise on *Meloidogyne*. Vol. II: Methodology. North Carolina State University. USA. Pp. 223.
- Dalmasso, A. (1969). Etude anatomique et taxonomique des genres *Xiphinema*, *Longidorus* et *Paralongidorus* (Nematoda: Dorylaimidae). *Memoires du Muséum National d'Histoire Naturelle. Paris, Serie A. Zoologie*, 61:33-82.
- Decraemer, W. and E. Geraet. (2006). Ectoparasitic nematodes. In: *Plant Nematology*. Edited by: R. Perry and M. Moens. CABI international. London, UK. p. 447.
- Jacob, J and J. van Bezooijen. (1984). *Manual for Practical Work in Nematology*. Wageningen. Pp. 77.
- Jaffee, B., A. Nyczepir and A. Golden. (1987). *Criconemella* spp. in ennsylvania peach orchards with morphological observations of *C. curvata* and *C. ornata*. *Journal of Nematology* 19:420-423.
- Knight, K., C. Barber and G. Page. (1997). Plant-parasitic nematodes of New Zealand recorded by host association. *Journal of Nematology*. 29(4):640-656.
- Lamberti, F. (1984). Nematode problems of the Mediterranean coastal strip in the Syrian Arab Republic. *Nematologia Mediterranea*, 12:53-64.
- Lamberti, F. and T. Bleve-Zacheo. (1979). Studies on *Xiphinema americanum* *sesu lata* with description of fifteen new species (Nematoda: Longidoridae). *Nematologia Mediterranea*. 7:51-106.

- McKenry, M. (1989). Nematodes. In: Peaches, Plums and Nectarines: Growing and Handling for Fresh Market. Editors: J. LaRue and R. Johnson, University of California. California. USA. Pp. 139-147.
- Meyer, A., and H. Hugo. (1994). *Xiphinema americanum* damaging peach trees in South Africa. *Journal of Nematology* 26:111 (Abstract).
- Nyczepir, A., and J. Halbrendt. (1993). Nematode pests of deciduous fruit and nut trees. pp. 381-425. In: Plant Parasitic Nematodes in Temperate Agriculture. Editors: K. Evans, D. Trudgill, and J. Webster. CAB International, Wallingford, England.
- Ogawa, J., E. Zehr, G. Bird, D. Richie, K. Uriu, and J. Uyemto. (1995). Compendium of Stone Fruit diseases. The American Phytopathological Society. p. 98.
- Sasser, J. and C. Carter. (1985). Overview of the international Meloidogyne project 1975-1984. In: Advanced Treatise on Meloidogyne. V1: Biology and Control. Edited by: J. Sasser, & C. Carter. North Carolina State University. Pp. 19-24.
- Sasser, J. and D. Freckman. (1987). A world perspective on nematology: The role of society. In: Vistas on Nematology. Edited by: J. Veech and A. Dickson. Society of Nematologists. USA. Pp. 7-14.
- Seinhorst, J. (1959). A rapid method for the transfer of nematodes from fixation to anhydrous glycerin. *Nematologica*, 4:67-69.
- Stirling, G. R. (1991). Biological Control of Plant Parasitic Nematodes, Progress, Problems and Prospects. CAB International. p. 282.
- Whitehead, A. G. (1998). Plant nematode control. CAB International. p. 384.

Received	2009/07/31	
Accepted for Publ.	2010/03/30	