
Misoprostol

*

Misoprostol

.2005

2 2004

2

:

:

18 :

:

(5)

1000

Misoprostol

Misoprostol

:

Misoprostol

-

-

-

*

The Place of Misoprostol in Management of Post Partum Hemorrhage (P.P.H) due to Uterine Atony

Tammam Al Ashkar *

Abstract

This study was organized to find out the role and the efficacy of Misoprostol in treating postpartum hemorrhage unresponsive to traditional agents that contract the uterus .

Place : Maternity hospital at Faculty of Medicine in Damascus university

Time : from 2, January, 2004 to 2, January, 2005 .

Number of patients : 18 patients who met the criteria in that period of time .

Criteria of selection : any patient with post partum hemorrhage due to uterine atony that did not respond well to the administration of Oxytocin and Metergin .

Misoprostol is an effective treatment of post partum hemorrhage due to uterine atony unresponsive to Oxytocin and Metergin.

Our study confirms the results of international studies of effectiveness of the use of misoprostol in treating post partum hemorrhage due to uterine atony .

Key Words : Misoprostol in Post Partum Hemorrhage Due to Uterine Atony

* Ass. Professor- Department of Obstetrics & Gynecology - Faculty of Medicine - Damascus University.

500 : **Introduction**
1000 Post Partum Hemorrhage

%10 ARDS

Causes of Post Partum Hemorrhage : **Definition**
Lacerations -1)
- -)
(8 1) : ()
Uterine Atony -2)
(20 1))
Coagulation -3 / (...)
) Disorders)
(...)
(...)
-4
Consumptive Coagulopathy

Misoprostol

		-2)
			.(
			-5
		-3	.Placenta Accreta
	:	Acute	-6
Oxytocin		-	.Inversion of the Uterus
	40-10	Rupture of the	-7
			.Uterus
0.2	Metergin	-	-8
			.Rupture of the Uterine scar
			-9
		-	
) Carboprost	Tromethamine		
250	(Hemabate		-10
90-15			
2			
800 Misoprostol		-	Management
	1000		: of Uterine Atony

(3) .

:Misoprostol

E1

(3) .

-
- :
-1
-2
-3
-4
-5

Misoprostol

3

Misoprostol

18 Misoprostol

Misoprostol

1000

Metergin

40-20

Oxytocin

18

Misoprostol

0.4

18

2 2004

2005

5-3

15 Misoprostol

Misoprostol

0.4 :

Metergin Misoprostol

1000

Misoprostol

15

18

Oxytocin

5-3

40-20

:

4	7	4	
5	4	3	
4	4	4	

% 83.3

%16.7

Misoprostol

%100	18	
%83.3	15	
%16.7	3	Misoprostol

Misoprostol

1000

14

Misoprostol

3

4

:

(5) 1998

18

15

3

3	%100	(4)
4	%83.3	

Misoprostol

:

1000

Misoprostol

(5)

Misoprostol

Metergin

Oxytocin

15

18

المراجع

1. Andolina, K, Daly, S, Roberts, N, et al , objective measurement of blood loss at delivery. Amj Obstet Gynecol 119,180,69.
2. Veland , K, Maternal Cardiovesical dynamics VII intrapartum Blood Volume changes , Amj Obstet Gynecol 1976, 126, 671.
3. Lokuganage, AV, Sullinan , KR , Niculescu , I , et al , a randomized study comparing redally administration of Misoprostol versus systometrine combined with an oxytocin infusion for the cession of primary postpartum hemorrhage , obstet Gynecol 2001, 80, 835 .
4. PCHO and SWNAGU , Mifeprystone and Misoprostol , Current Obestetrics and gynecology , 1999 , 9 , 29 , 33 .
5. O'Brien P , El Refary , H , Gordon , A , et al , Rectally Administration of Misoprostol for the treatment of postpartum hemorrhage unresponsive to Oxytocin and ergotamine , a descriptive study , Obestet Gynecol , 1998 , 92 , 212

.2006/2/8:

.2006/4/13: